

# Reducing the Length of Mechanical Ventilation with Significance

## A Case Study of Sample Size Estimation Trial Design using Monte-Carlo Simulation

Yeong Shiong Chiew<sup>1</sup>, Christopher Pretty<sup>1</sup>, Elena Molchanova<sup>2</sup>, Carl Scarrott<sup>2</sup>,  
Daniel Redmond<sup>1</sup>, Geoffrey M Shaw<sup>3</sup>, J. Geoffrey Chase<sup>1</sup>

<sup>1</sup> Department of Mechanical Engineering, University of Canterbury,  
Christchurch, New Zealand

<sup>2</sup> School of Mathematics and Statistics, University of Canterbury,  
Christchurch, New Zealand

<sup>3</sup> Intensive Care Unit, Christchurch Hospital,  
Christchurch, New Zealand

# Mechanical Ventilation

- Primary support for patients with respiratory failure
  - Maintaining lung recruitment
  - Effective O<sub>2</sub> delivery
  - Provide good patient-ventilator interaction without inducing further stress



# Mechanical Ventilation

- Primary support for patients with respiratory failure
  - Maintaining lung recruitment
  - Effective O<sub>2</sub> delivery
  - Provide good patient-ventilator interaction without inducing further stress
- What is considered as effective ventilation?
  - No further injuries
  - Patient receiving sufficient O<sub>2</sub>
  - Good patient interaction with ventilation
  - Recovery



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?

# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH, PCO<sub>2</sub>, PaO<sub>2</sub>



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH,  $\text{PCO}_2$ ,  $\text{PaO}_2$
  - $\text{SPO}_2$  as desaturation events: drop less than 88%



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH,  $\text{PCO}_2$ ,  $\text{PaO}_2$
  - $\text{SPO}_2$  as desaturation events: drop less than 88%
  - Asynchrony index




# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH, PCO<sub>2</sub>, PaO<sub>2</sub>
  - SPO<sub>2</sub> as desaturation events: drop less than 88%
  - Asynchrony index
  - Multiple organ dysfunction (MOD)

## Onset of Multiple Organ Dysfunction Syndrome (MODS)

IAP > 20 mmHg





# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH,  $\text{PCO}_2$ ,  $\text{PaO}_2$
  - $\text{SPO}_2$  as desaturation events: drop less than 88%
  - Asynchrony index
  - Multiple organ dysfunction (MOD)
  - Respiratory or other haemodynamic parameters



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH,  $\text{PCO}_2$ ,  $\text{PaO}_2$
  - $\text{SPO}_2$  as desaturation events: drop less than 88%
  - Asynchrony index
  - Multiple organ dysfunction (MOD)
  - Respiratory or other haemodynamic parameters
  - Ventilator dependency: Weaning process



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH, PCO<sub>2</sub>, PaO<sub>2</sub>
  - SPO<sub>2</sub> as desaturation events: drop less than 88%
  - Asynchrony index
  - Multiple organ dysfunction (MOD)
  - Respiratory or other haemodynamic parameters
  - Ventilator dependency: Weaning process
  - Length of mechanical ventilation or ventilator free days



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH, PCO<sub>2</sub>, PaO<sub>2</sub>
  - SPO<sub>2</sub> as desaturation events: drop less than 88%
  - Asynchrony index
  - Multiple organ dysfunction (MOD)
  - Respiratory or other haemodynamic parameters
  - Ventilator dependency: Weaning process
  - Length of mechanical ventilation or ventilator free days
  - Mortality





# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- What do we measure?
  - Arterial blood gasses: pH, PCO<sub>2</sub>, PaO<sub>2</sub>
  - SPO<sub>2</sub> as desaturation events: drop less than 88%
  - Asynchrony index
  - Multiple organ dysfunction (MOD)
  - Respiratory or other haemodynamic parameters
  - Ventilator dependency: Weaning process
  - **Length of mechanical ventilation or ventilator free days**
  - Mortality



# How do we measure effective ventilation?

- What is the surrogate of an effective mechanical ventilation?
- Why Ventilator Free Days (VDF) or Length of Mechanical ventilation (LOMV)?



Cooper LM, Linde-Zwirble WT: **Medicare intensive care unit use: Analysis of incidence, cost, and payment.** *Critical Care Medicine* 2004, **32**:2247-2253.

Dasta JF, McLaughlin TP, Mody SH, Piech CT: **Daily cost of an intensive care unit day: The contribution of mechanical ventilation.** *Crit Care Med* 2005, **33**:1266-1271.

Zilberberg M, Shorr A: **Prolonged acute mechanical ventilation and hospital bed utilization in 2020 in the United States: implications for budgets, plant and personnel planning.** *BMC Health Services Research* 2008, **8**:242.

# Existing studies with LoMV as primary or secondary outcome

Study	No. Patient	Outcome	Groups (Number of patient) LoMV or Vent Free Days (in mean $\pm$ standard deviation or median [interquartile range])		p-value
ARDSNet (The Acute Respiratory Distress Syndrome Network, 2000)	861	VFD*	Low Vt+ (432) 12 $\pm$ 11	High Vt (429) 10 $\pm$ 11	0.0070
ALVEOLI (Brower et al., 2004)	549	VFD	Lower PEEP# (273) 14.5 $\pm$ 10.4	Higher PEEP (276) 13.8 $\pm$ 10.6	0.5000
EXPRESS (Mercat et al., 2008)	767	VFD	Minimal distension (382) 3 [0-17]	Increased recruitment (385) 7 [0-19]	0.0400
LOVS (Meade et al., 2008)	983	LoMV	Control (507) 10 [6-16]	Lung open (475) 10 [6-17]	0.9200
Meta-Analysis (Briel et al., 2010)	2299	VFD	Lower PEEP (1136) 11 [0-21]	Higher PEEP (1163) 13 [0-22]	0.1000
Individualised PEEP (Pintado et al., 2013)	70	VFD	Control (36) 0 [0-15.75]	Intervention (34) 1 [0-18]	0.1600
Sedation Study (Strøm et al., 2010)	113	VFD	Control (58) 18.0 [0-24.1]	No Sedation (55) 6.9 [0-20.5]	0.0191

# Existing studies with LoMV as primary or secondary outcome

Study	No. Patient	Outcome	Groups (Number of patient) LoMV or Vent Free Days (in mean $\pm$ standard deviation or median [interquartile range])		p-value
ARDSNet (The Acute Respiratory Distress Syndrome Network, 2000)	861	VFD*	Low Vt+ (432) 12 $\pm$ 11	High Vt (429) 10 $\pm$ 11	0.0070
ALVEOLI (Brower et al., 2004)	549	VFD	Lower PEEP# (273) 14.5 $\pm$ 10.4	Higher PEEP (276) 13.8 $\pm$ 10.6	0.5000
EXPRESS (Mercat et al., 2008)	767	VFD	Minimal distension (382) 3 [0-17]	Increased recruitment (385) 7 [0-19]	0.0400
LOVS (Meade et al., 2008)	983	LoMV	Control (507) 10 [6-16]	Lung open (475) 10 [6-17]	0.9200
Meta-Analysis (Briel et al., 2010)	2299	VFD	Lower PEEP (1136) 11 [0-21]	Higher PEEP (1163) 13 [0-22]	0.1000
Individualised PEEP (Pintado et al., 2013)	70	VFD	Control (36) 0 [0-15.75]	Intervention (34) 1 [0-18]	0.1600
Sedation Study (Strøm et al., 2010)	113	VFD	Control (58) 18.0 [0-24.1]	No Sedation (55) 6.9 [0-20.5]	0.0191



## **This study aimed to:**


- Estimate the sample size required for a randomised controlled trial for mechanical ventilation.
  - Sample size for the change in LoMV detection
- We compare two methods
  - Using model-based method (Altman-nomogram)
  - Using simulation-based method (Monte-Carlo)
- Using ICU patients data from a single hospital Intensive care unit (From 2011-2013)

# Christchurch ICU Record

- Age, Apache diagnostic code, ICU mortality, length of mechanical ventilation, ventilation modes.
- Of the 3907 patients admitted, 2921 patients (75%) required MV, and 2534 (65%) patients were invasively ventilated

# Christchurch ICU Record

- Age, Apache diagnostic code, ICU mortality, length of mechanical ventilation, ventilation modes.
- Of the 3907 patients admitted, 2921 patients (75%) required MV, and 2534 (65%) patients were invasively ventilated




- Cohort A LoMV mean ( $\pm$  SD) LoMV for Cohort A is  $3.23 \pm 7.03$  days (median = 0.72 days [IQR: 0.24-2.62]).

# Altman Nomogram


# Altman Nomogram

Standardised  
Difference  
=  
 $\Delta\text{mean} / \text{SD}$




# Altman Nomogram

3 day difference  
Standardised  
Difference




Power

# Altman Nomogram




# Altman Nomogram






# Altman Nomogram



# Altman Nomogram



# Monte-Carlo Simulation

- Altman Nomogram assumes normally distributed data sets, and thus, not suitable for data that are skewed to the left.
- Hence, we tried a simulation based approach to estimated sample size

# Monte-Carlo Simulation

- Altman Nomogram assumes normally distributed data sets, and thus, not suitable for data that are skewed to the left.
- Hence, we tried a simulation based approach to estimated sample size
- Using Simulation based approach, we are able to impose a selection criteria on patients
  - (1) Patients who are likely to be discontinued from MV within 24 hours
  - (2) Patients with increase intracranial pressure
  - (3) Patients who have significant weakness from any neurological disease
  - (4) Patients who have asthma as the primary presenting condition or a history of significant chronic obstructive pulmonary disease, and
  - (5) Patients who are pregnant
- How do we know which patient? APACHE II Diagnostic codes

# Christchurch ICU Record

	Code	Description		Code	Description
Non - Operative	100	Cardiovascular	Post - Operative	1200	Cardiovascular
	200	Respiratory		1300	Respiratory
	300	Gastrointestinal		1400	Gastrointestinal
	400	Neurological		1500	Neurological
	500	Sepsis		1600	Trauma
	600	Trauma		1700	Renal/ Genitourinary
	700	Metabolic		1800	Gynaecological
	800	Haematology		1900	Musculoskeletal
	900	Renal disorder		2100	Haematological
	1000	Other medical disorders		2200	Metabolic
	1100	Musculoskeletal/ Skin disease			
	o	No diagnosis entered			

# Monte-Carlo Simulation

- Non-operative neurological (400)
- Post-operative neurological (1500)
- Chronic obstructive pulmonary disease (206)
- Asthma (209)
- Head trauma with or without multi trauma (601)
- Multi trauma with spinal injury (604)
- Isolated cervical spine injury (605)
- Post operation patients: head trauma with or without multi trauma (1601)
- Post operation patients: multi trauma with spinal injury (1604)
- Post operation patients: isolated cervical spine injury (1605)
- Pregnancy-related disorder (1802).
- Patients with LoMV less than 1 day and more than 30 days were also excluded.



These exclusion criteria were chosen based the clinical implication that these patient may not benefit from a MV intervention, or could be harmed in some cases.

# Monte-Carlo Simulation

- After imposing the exclusion criteria, the number of patients who might benefit and be eligible for the study was reduced to 744 (19% of total patients admitted to ICU or 29% of patients requiring invasive MV).

# Monte-Carlo Simulation



- After imposing the exclusion criteria, the number of patients who might benefit and be eligible for the study was reduced to 744 (19% of total patients admitted to ICU or 29% of patients requiring invasive MV).
- Cohort B mean LoMV was  $5.81 \pm 6.30$  days (median = 2.92 days [IQR: 1.67-7.38]),





# Monte-Carlo Simulation

- After imposing the exclusion criteria, the number of patients who might benefit and be eligible for the study was reduced to 744 (19% of total patients admitted to ICU or 29% of patients requiring invasive MV).
- Cohort B mean LoMV was  $5.81 \pm 6.30$  days (median = 2.92 days [IQR: 1.67-7.38]),
- Significantly different from Cohort A ( $p < 0.05$  using Student *t*-test and Wilcoxon ranksum test and Kolmogorov-Smirnov test).



# Monte-Carlo Simulation

	Step	Description	Tests
1.	Patient Cohorts	<ul style="list-style-type: none"><li>Select a patient cohort:<ol style="list-style-type: none"><li>Cohort A includes all invasively ventilated patients.</li><li>Cohort B is created from Cohort A by imposing exclusion criteria.</li></ol></li></ul>	<ul style="list-style-type: none"><li>Cohort A</li><li>Cohort B</li></ul>

# Monte-Carlo Simulation

	Step	Description	Tests
1.	Patient Cohorts	<ul style="list-style-type: none"> <li>Select a patient cohort:                             <ol style="list-style-type: none"> <li>Cohort A includes all invasively ventilated patients.</li> <li>Cohort B is created from Cohort A by imposing exclusion criteria.</li> </ol> </li> </ul>	<ul style="list-style-type: none"> <li>Cohort A</li> <li>Cohort B</li> </ul>
2.	Sample Size Selection	<ul style="list-style-type: none"> <li>Randomly select patients from the patient cohort and assign each patients to a treatment group. 1) Control group or 2) Intervention group.</li> <li>Patient is selected with replacement.</li> <li>Various sample size of each treatment group is tested.</li> </ul>	<ul style="list-style-type: none"> <li>Total sample size N= 200, 300, 400... 5000 patients.</li> <li>Or sample size with higher resolution 200, 220, 240... 600 patients</li> </ul>

# Monte-Carlo Simulation

	Step	Description	Tests
1.	Patient Cohorts	<ul style="list-style-type: none"> <li>Select a patient cohort:                             <ol style="list-style-type: none"> <li>Cohort A includes all invasively ventilated patients.</li> <li>Cohort B is created from Cohort A by imposing exclusion criteria.</li> </ol> </li> </ul>	<ul style="list-style-type: none"> <li>Cohort A</li> <li>Cohort B</li> </ul>
2.	Sample Size Selection	<ul style="list-style-type: none"> <li>Randomly select patients from the patient cohort and assign each patients to a treatment group. 1) Control group or 2) Intervention group.</li> <li>Patient is selected with replacement.</li> <li>Various sample size of each treatment group is tested.</li> </ul>	<ul style="list-style-type: none"> <li>Total sample size N= 200, 300, 400... 5000 patients.</li> <li>Or sample size with higher resolution 200, 220, 240... 600 patients</li> </ul>
3.	Difference in LoMV	<ul style="list-style-type: none"> <li>Impose a difference in LoMV between two groups.</li> <li>The LoMV in Intervention group is reduced by the chosen percentage.</li> </ul> $\text{LoMV intervention} = \text{LoMV patient} \times (100\% - \text{Percentage reduction})$ <ul style="list-style-type: none"> <li>The difference in LoMV ranges from 5 to 30% of total LoMV.</li> </ul>	<ul style="list-style-type: none"> <li>Difference of LoMV = 5, 10, 15, ... 30%</li> </ul>


# Monte-Carlo Simulation

	Step	Description	Tests
1.	Patient Cohorts	<ul style="list-style-type: none"> <li>Select a patient cohort:                             <ol style="list-style-type: none"> <li>Cohort A includes all invasively ventilated patients.</li> <li>Cohort B is created from Cohort A by imposing exclusion criteria.</li> </ol> </li> </ul>	<ul style="list-style-type: none"> <li>Cohort A</li> <li>Cohort B</li> </ul>
2.	Sample Size Selection	<ul style="list-style-type: none"> <li>Randomly select patients from the patient cohort and assign each patients to a treatment group. 1) Control group or 2) Intervention group.</li> <li>Patient is selected with replacement.</li> <li>Various sample size of each treatment group is tested.</li> </ul>	<ul style="list-style-type: none"> <li>Total sample size N= 200, 300, 400... 5000 patients.</li> <li>Or sample size with higher resolution 200, 220, 240... 600 patients</li> </ul>
3.	Difference in LoMV	<ul style="list-style-type: none"> <li>Impose a difference in LoMV between two groups.</li> <li>The LoMV in Intervention group is reduced by the chosen percentage.  <math display="block">\text{LoMV intervention} = \text{LoMV patient} \times (100\% - \text{Percentage reduction})</math> </li> <li>The difference in LoMV ranges from 5 to 30% of total LoMV.</li> </ul>	<ul style="list-style-type: none"> <li>Difference of LoMV = 5, 10, 15, ... 30%</li> </ul>
4.	Statistical Test	<ul style="list-style-type: none"> <li>Perform statistical test comparing the LoMV between two groups.</li> <li>Using parametric and non-parametric tests.</li> <li>A value of <math>p &lt; 0.05</math> indicates that LoMV for intervention group is significantly different from control group.</li> </ul>	<ul style="list-style-type: none"> <li>Student t-test</li> <li>Student t-test (log scale)</li> <li>Wilcoxon Ranksum test</li> <li>Kolmogorov-Smirnov test</li> </ul>

# Monte-Carlo Simulation


	Step	Description	Tests
1.	Patient Cohorts	<ul style="list-style-type: none"> <li>Select a patient cohort:                             <ol style="list-style-type: none"> <li>Cohort A includes all invasively ventilated patients.</li> <li>Cohort B is created from Cohort A by imposing exclusion criteria.</li> </ol> </li> </ul>	<ul style="list-style-type: none"> <li>Cohort A</li> <li>Cohort B</li> </ul>
2.	Sample Size Selection	<ul style="list-style-type: none"> <li>Randomly select patients from the patient cohort and assign each patients to a treatment group. 1) Control group or 2) Intervention group.</li> <li>Patient is selected with replacement.</li> <li>Various sample size of each treatment group is tested.</li> </ul>	<ul style="list-style-type: none"> <li>Total sample size N= 200, 300, 400... 5000 patients.</li> <li>Or sample size with higher resolution 200, 220, 240... 600 patients</li> </ul>
3.	Difference in LoMV	<ul style="list-style-type: none"> <li>Impose a difference in LoMV between two groups.</li> <li>The LoMV in Intervention group is reduced by the chosen percentage.  <math display="block">\text{LoMV intervention} = \text{LoMV patient} \times (100\% - \text{Percentage reduction})</math> </li> <li>The difference in LoMV ranges from 5 to 30% of total LoMV.</li> </ul>	<ul style="list-style-type: none"> <li>Difference of LoMV = 5, 10, 15, ... 30%</li> </ul>
4.	Statistical Test	<ul style="list-style-type: none"> <li>Perform statistical test comparing the LoMV between two groups.</li> <li>Using parametric and non-parametric tests.</li> <li>A value of <math>p &lt; 0.05</math> indicates that LoMV for intervention group is significantly different from control group.</li> </ul>	<ul style="list-style-type: none"> <li>Student t-test</li> <li>Student t-test (log scale)</li> <li>Wilcoxon Ranksum test</li> <li>Kolmogorov-Smirnov test</li> </ul>
5.	Power Analysis	<ul style="list-style-type: none"> <li>Each Monte-Carlo simulation iteration will generate a p-value for each statistical test.</li> <li>For a given sample size and significance level <math>\alpha</math>, statistical power is evaluated as the proportion of iterations for which the <math>p &lt; \alpha</math>.</li> </ul>	<ul style="list-style-type: none"> <li>E.g. for 10000 Monte-Carlo iterations, if <math>p &lt; \alpha</math> for 84% (8400 iterations), Power = 0.84.</li> </ul>

# Monte-Carlo Simulation




# Monte-Carlo Simulation


5% LoMV Difference




10% LoMV Difference




15% LoMV Difference




20% LoMV Difference



25% LoMV Difference




30% LoMV Difference






# Monte-Carlo Simulation


5% LoMV Difference




10% LoMV Difference




15% LoMV Difference




20% LoMV Difference



25% LoMV Difference



30% LoMV Difference



# Discussion


- One of the ideal inclusion criteria in other studies is to focus on patients with a more severe form of respiratory failure, such as the acute respiratory distress syndrome (ARDS).
- LoMV distributions may vary between centres ([Van Der Lee et al., 2009](#)). This variability in patient distribution means that the  $N_{\text{total}}$  derived from this study may only be applicable to the participating centre or other regional centres that have similar characteristics ([Van Der Lee et al., 2009](#)).
- The changes of LoMV used in this study were arbitrarily chosen and may not represent the true possible LoMV change for any given intervention ([Schulz and Grimes, 2005](#)).
- Thus, given the effective sample size and percentage difference in LoMV needed, the results suggest that the clinical outcome (LoMV) requires a large sample size due to the high variability in the population.

# Final Thoughts

- Trial inclusion and exclusion criteria is important to capture the targeted group.
  - A specific group or,
  - A generalised group
- There are other co-founding factors that can affect MV deliveries.
  - Intention to treat
  - Protocol deviation
  - Effective protocol
  - MV is a supporting therapy

# Acknowledgement

- New Zealand Health Research Council
- Centre of Bioengineering (University of Canterbury, New Zealand)
- GIGA Cardiovascular Science (University of Liege, Belgium)
- Institute of Technical Medicine (Furtwangen University, Germany)
- Department of Control Engineering and Information Technology (Budapest University of Technology and Economics, Hungary)



# Acknowledgement

*This work was supported by EU FP7 IRSES (FP7-PEOPLE-2012-IRSES) program, project title: eTime - Engineering Technology-based Innovation in Medicine, Grant No. 318943.*





תודה  
 Dankie Gracias شكراً  
 Спасибо Merci Takk  
 Köszönjük Terima kasih  
 Grazie Dziękujemy Děkojame  
 Ďakujeme Vielen Dank Paldies  
 Kiitos Täname teid 谢谢  
**Thank You** Tak  
 感谢您 Obrigado Teşekkür Ederiz  
 Σας Ευχαριστούμ 감사합니다  
 ඔබට  
 Bedankt Děkuje vám  
 ありがとうございます  
 Tack

